

GENERAL INDEX TO VOLUME XXV

New scientific names of plants and the final members of new combinations are printed in **bold face type**; synonyms and page numbers having reference to figures and plates, in *italics*; and previously published names and all other matter, in ordinary type.

A

- Abacina*, 550
 Acarosporaceae, 554
 Acid reaction of pollen granules of
 Tradescantia, 460
Actinodaphne, 400; *angustifolia*, 418;
chinensis, 386, 414, var. *oblongifolia*,
 386, 414, var. *rotundifolia*, 385, 414;
cochinchinensis, 404; confertiflora,
 400, 401; *confertifolia*, 414, 416;
cupularis, 399, 405; ferruginea, 407;
glauca, 410; Henryi, 407; *hong-*
kongensis, 414, 418; Hookeri, 414;
lanceifolia, 406, var. *sinensis*, 406;
Lecomtei, 399, 413; *litseaefolia*, 408;
magniflora, 403; obovata, 399, 404;
omeiensis, 411; pedunculata, 414;
perluca, 409; pilosa, 404; reticu-
 lata, 412, var. *Forrestii*, 412, var.
glabra, 412; *reticulata*, 413, var.
omeiensis, 411; *rotundifolia*, 385,
 414; sesquipedalis, 409, var. *cam-*
bodiana, 409; *setchenensis*, 411;
trichocarpa, 402; Tsaii, 408
 Additions, corrections, and revisions to
 the "Annotated Catalogue of the
 Flowering Plants of Missouri," 775
Adinandra lasiopetala, 355
Agalopisma, 622
Alectoria, 598; **antarctica**, 599, 600
 Algae in Marie Byrd Land, 467, 495,
 520
 Algal components of lichens, 520, 735;
 of *Stereocaulon*, 739, 741, 743, 750
 Alkali reaction of microspore granules
 of *Tradescantia*, 460
 Allen, Caroline K. Studies in the Lau-
 raceae. I. Chinese and Indo-Chinese
 species of *Litsea*, *Neolitsea*, and *Ac-*
tinodaphne, 361
Alonsoa, 438; *acutifolia*, 453; *caulia-*
lata, 453; *incisifolia*, 453; *linearifolia*,
 453; *linearis*, 453; *Mutisii*, 453; *pro-*
cumbens, 453
Alsine chamaedrios, 446
Alsinas seu Spergulae dictae species
africana, 444
Amarylhidaceae, Panamanian, 824

ANN. MO. BOT. GARD., VOL. 25, 1938.

- Amelanchier canadensis f. nuda*, 772
Amphiloma, 622
Anagallis capensis, 446, 452; *foliis sinu-*
atis, 437; *purpurea*, 446
Anisophyllum (sect.) 2, 10
Anisophyllum, 2, 10; *cordatum*, 62;
multiforme, 45; *nodosum*, 10; *Thymi-*
folium, 67; *virgatum*, 59
 Antarctic Expedition, The second Byrd
 —Botany, 467
Antarctica, botanical exploration of,
 516, 517
Antirrhinum, 435
Anzia sect. *Pannoparmelia*, 586
 Apocynaceae: Panamanian, 832;
 Studies in the. VII. An evaluation
 of the genera *Plumeria* L. and *Hima-*
anthus Willd., 189
 Apothecial characters of *Stereocaulon*,
 correlation of, with cephalodial types,
 746
Ardisia furfuracella, 832
Argopsis, 734, 736
Asclepiadaceae, Panamanian, 833
Aspidosperma, 194
Audibertia (sect.), 98
Audibertia, 98; *capitata*, 123; *Cleve-*
landii, 119; *decurrens*, 110; *Dorrii*,
 133; *grandiflora*, 110; *humilis*, 136;
incana, 130, var. *pachystachya*, 125,
 var. *pilosa*, 133; *Mearnsii*, 131;
nivea, 121; *pachystachya*, 125;
Palmeri, 146; *polystachya*, 140; *spin-*
ulosa, 113; *stachyoides*, 112, var.
revoluta, 117; *Vaseyi*, 138
Audibertiella, 98; *argentea*, 131; *cap-*
itata, 123; *Clevelandii*, 119; *Dorrii*,
 133; *grandiflora*, 110; *humilis*, 136;
incana, 130; *nivea*, 121; *Palmeri*,
 146; *polystachya*, 140; *stachyoides*,
 113; *Vaseyi*, 138

B

- Baeomyces, 734
 Baker, Gladys E., Carroll W. Dodge
 and. The second Byrd Antarctic Ex-
 pedition—Botany. II. Lichens and
 lichen parasites, 515

- Barbula Byrdii*, 719, 720
 Bartram, Edwin B. The second Byrd Antarctic Expedition—Botany. III. Mosses, 719
Begonia Allenii, 830
Begoniaceae, Panamanian, determined by P. C. Standley, 830
Benzoïn fruticosum, 399; *Levinci*, 423; *obovatum*, 367; *puberulum*, 370; *setchuensis*, 411
Berengeria, 655
Biatorella arachnoidea, 555, 680
Biatoridium, 554
 Birds, relation of, to Antarctic plant life, 485, 501, 502, 504
Blastenia, 611; *grisea*, 613, 680, 694; subg. *Protoblastenia*, 605; *succinea*, 612, 694
Blasteniaceae, 604
Botrytis, 522, 712
Bryaceae, 723
Bryum (*Argyrobryum*) *antarcticum*, 719, 724; *austropolare*, 724; *Siplei*, 723
Buchenavia capitata, 831
Buellia, 631; *albida*, 654, 708; *albordiana*, 647, 700, 702; *brunnescens*, 646, 700; sect. *Catocarpus*, 551; *chrysea*, 642, 700; *dendritica*, 651, 706, 708; *flavoplana*, 640, 698, 718; *floccosa*, 650, 702, 704; *frigida*, 653, 702, 718; *grisea*, 639, 706; *musciola*, 643, 706; *olivaceobrunnea*, 636, 704; *pallida*, 638, 700; *Russellii*, 649, 708; *Siplei*, 635, 708, 710; *stellata*, 644, 698, 700
Buelliaceae, 521, 631
Burmanniaceae, Panamanian, determined by F. P. Jonker, 825
 Byrd Antarctic Expedition, The second—Botany. I. Ecology and geographical distribution, 467; II. Lichens and lichen parasites, 515; III. Mosses, 719
- C
- Cacalia arborea*, 798; *pulchella*, 816; *teretifolia*, 819
 Californian *Salvias*, The, 95
Callopisma, 619
Caloplaca, 619; *athallina*, 621; *aurantiaca*, 622; *cinericola*, 622; *cirrochrooides*, 626; *inordinata*, 626; *Joannae*, 626; *lucens*, 627; *sublobulata*, 627
Candelariella, 580; *albovirens*, 582, 684, 686; *chrysea*, 580, 686
 Carbohydrates, tests for, in microspore granules of *Tradescantia*, 458
Cardamine bulbosa f. *fontinalis*, 771
Carya Buckleyi var. *arkansana* f. *glabra*, 770
 "Catalogue, Annotated, of the Flowering Plants of Missouri," Additions, corrections, and revisions to the, 775
Catillaria, 543; *arachnoidea*, 549, 672, 718; *cremea*, 544, 672; *floccosa*, 545, 672; *granulosa*, 547, 674; *inconspicua*, 548, 674
Catocarpon, 551
Catocarpus, 551
Cephalodia of *Stereocaulon*: historical review of, 729; illustrating their varied gross morphology, 764; origin and development of, 738, 766, 768; physiological significance of, 737; The taxonomic importance and phylogenetic significance of the, 729; types of, 737, 745, correlation of apothecial characters with, 746, of vegetative characters with, 748, of geographical distribution with, 758, botryose, 741, serobiculate, 742, spherical, 739
Chamaesyce, 2; *atrococca*, 42; *Celastroides*, 24; *Clusiaefolia*, 10; *Degenert*, 62; *festiva*, 61; *Forbesii*, 12; *halemanui*, 23; *Hillebrandii*, 56, var. *waimanoana*, 58; *Hookeri*, 59; *Hypericifolia*, 66; *lorifolia*, 33; *multiformis*, 45; *olowaluana*, 40; *Remyi*, 15; *Rockii*, 14; *Skottsbergii*, 54
 Chemical studies, Optical and, on the granules in microspores of *Tradescantia*, 455, 458
 Chester Mts., geology and plant life of, 493, 514
 Chinese and Indo-Chinese species of *Litsea*, *Neolitsea*, and *Actinodaphne*, 361
Chiococca phaenostemon, 838
Chlorophyceae, 520, 751
Cladonia, 734, 745
Cladoniaceae, phylogenetic relationships in, 733, 735
 Claude Swanson Mts., 487, 506
Cleyera, 299, 300
Cogswellia, 225, 228; *alata*, 287; *altensis*, 288; *ambigua*, 247; *angustata*, 268; *anomala*, 285; *argensis*, 263; *artemisiarum*, 264; *Austinae*, 257; *bicolor*, 246; *Brandegei*, 290; *brecciarum*, 287; *brevifolia*, 286; *Canbyi*, 244; *caruifolia*, 252, var. *patens*, 250, var. *Vaseyi*, 251; *Chandleri*, 250; *circumdata*, 249; *concinna*, 276; *Congdoni*, 266; *Cottami*, 277; *Cous*, 247; *Cusickii*, 287; *dasycarpa*, 262; *daucifolia*, 260; *decipiens*, 255;

- Donnellii*, 271; *duchesnensis*, 292; *Eastwoodae*, 275; *elliptica*, 265; *farinosa*, 242; *flava*, 264; *foeniculacea*, 259; *fragrans*, 271; *Geyeri*, 243; *gigantea*, 285; *Gormani*, 240, f. *purpurea*, 241; *Grayi*, 270; *Hallii*, 279; *Hasselti*, 236; *Hendersoni*, 244; *Howellii*, 237; *insularis*, 239; *Jaredi*, 262; *Jonesii*, 258; *juniperina*, 259; *laevigata*, 288; *lapidosa*, 292; *latifolia*, 291; *Leibergii*, 279; *Lemmoni*, 292; *leptocarpa*, 246; *leptophylla*, 283; *lucida*, 236; *MacDougalli*, 258; *macrocarpa*, 264, 279, var. *artemisiarum*, 264; *marginata*, 253; *Martindalei*, 280, var. *angustata*, 268; *microcarpa*, 279; *millefolia*, 270, var. *depauperata*, 270; *minima*, 273; *mojavensis*, 263; *montana*, 248; *nevadensis*, 254, var. *cupulata*, 254, var. *Parishii*, 255, var. *pseudorientalis*, 255; *nudicaulis*, 291; *Nuttallii*, 278, var. *alpina*, 279; *oregana*, 273; *orientalis*, 261; *orogenioides*, 242; *pallida*, 239; *Parishii*, 255; *Parryi*, 277; *parvifolia*, 238; *Piperi*, 241; *platycarpa*, 283; *platyphylla*, 291; *Plummerae*, 256; *rigida*, 240; *robustior*, 286; *Sandbergii*, 266; *scabra*, 275; *serpentina*, 271; *simplex*, 283; *simulans*, 264; *Sonnei*, 257; *Suksdorfii*, 289; *tomentosa*, 263; *Torreyi*, 267; *triternata*, 284, var. *alata*, 287, f. *lancifolia*, 285, var. *robustior*, 286; *utriculata*, 250, var. *papillata*, 251; *vaginata*, 249; *Vaseyi*, 251; *villosa*, 259; *Watsoni*, 245
- Collemiaceae, 520
- Combretaceae, Panamanian, determined by P. C. Standley, 831
- Coniocarpineae, 736
- Conopholis americana, 836; mexicana, 835, 836; panamensis, 835, 835
- Contributions toward a flora of Panama. II. Miscellaneous collections during 1936-1938, 823
- Corrections to the "Annotated Catalogue of the Flowering Plants of Missouri," 775
- Cranichis muscosa, 825
- Crinum darienensis, 824; erubescens, 825
- Culcitium magellanicum, 809
- Cyclocarpineae, 733
- Cynoches chlorochilon, 825
- Cylicodaphne longipes, 379
- Cynopteris duchesnensis, 292; longipes, 292; Newberryi, 292
- Cynomarathrum, 228; alpinum, 279; Brandegei, 290; brecciarum, 287; Eastwoodae, 275; latilobum, 281; Macbridei, 292; megarrhizum, 282; Nuttallii, 278; Parryi, 277; scabrum, 275
- D
- Daphnidium elongatum, 390; lancifolium, 406
- Diascia, 436, 438, 446; Bergiana, 436, 437, 438; diffusa, 435, 436, 437, 453; elongata, 453; macrophylla, 437, 440, 453; montana, 446; Scullyi, 446; unilabiata, 453
- Diastase, effect of, on microspore granules of Tradescantia, 459
- Dictyorinis, 655
- Didymodon glaciale, 719
- Dimelaena, 655
- Diospyros cerasifolia, 326; serrata, 321
- Diphais, 550
- Diphanis, 550
- Diplonaevia Parmelliae, 661, 712
- Dodge, Carroll W. and Gladys E. Baker. The Second Byrd Antarctic Expedition—Botany. II. Lichens and lichen parasites, 515
- Douglasiana, 101
- Dryopteris Hostmanni, 824; Lindigii, 824
- Durham Point, plant life of, 496
- E
- Echinospache (subsect.), 100
- Echinospache (sect.), 100
- Echites turbinata, 833
- Ecology and geographical distribution of the Antarctic flora, 467
- Edsel Ford ranges of Marie Byrd Land, 468; chemical analyses of some of the igneous mountains in, 474; general view of, 504; temperature of, 480
- Elaphoglossum siliquoides, 824
- Enzyme reactions of pollen granules of Tradescantia, 459
- Epling, Carl. The Californian Salvias. A review of Salvia, section Audibertia, 95
- Eueurya (sect.), 299, 306
- Eulophus ambiguus, 246; triternatus, 284
- Euphorbia L., Revision of the Hawaiian species of, 1
- Euphorbia, 5; Arnottiana, 59; atrococca, 42, var. kilaeana, 44, var. kokeana, 44; bifida, 66; Celastroides, 24, var. amplexens, 36, 38, var. halawana, 31, 34, f. kahanana, 31, var. hanapepensis, 35, 36, var. haupuana, 29, var. Humbertii, 32, var. kaenana,

- 28, 82, var. *kealiana*, 30, var. *kohaliana*, 30, var. *lorifolia*, 33, var. *mauiensis*, 38, var. *moomomiana*, 27, 80, var. *nematopoda*, 33, var. *niuensis*, 36, var. *saxicola*, 32, var. *Stokesii*, 26, var. *waikoluensis*, 29; *Celastroides*, 35; *Clusiaefolia*, 10; *Clusiaefolia* var. *grandifolia*, 13; *cordata*, 62; *coriariaefolia*, 59; *cyathophora*, 72; *Degeneri*, 62, var. *molokaiensis*, 64, 94; *festiva*, 61; *Forbesii*, 12; *geniculata*, 72; *halemanui*, 23, 78; *heterophylla*, 70, var. *cyathophora*, 72; *heterophylla* var. *genuina*, 70; *Hillebrandii*, 56, var. *palikeana*, 58, var. *waimanoana*, 58; *hirta*, 64; *Hookeri*, 59, var. *integrifolia*, 61; *Hypericifolia*, 66; *lorifolia*, 33, 36, 39, var. *gracilis*, 41; *multiformis*, 45, var. *haleakalana*, 49, var. *kaalana*, 52, var. *kapuleiensis*, 50, var. *manoana*, 50, var. *microphylla*, 47, var. *perdita*, 53, var. *sparsiflora*, 51, var. *tomentella*, 52, 90; *multiformis*, 47, 50, var. *a*, 47, var. *Celastroides*, 24, var. *lorifolia*, 33, var. *tenuior*, 47; *myrtifolia*, 59; *oahuensis*, 32; *olowaluana*, 40, var. *gracilis*, 41; *palustris*, 51; *Peplus*, 73; *pilulifera*, 64; *prostrata*, 68; *Remyi*, 15, var. *hanaleiensis*, 23, var. *kahili-ana* (*kalihi-ana*), 19, var. *kauaiensis*, 22, var. *leptopoda*, 18, var. *Lydgatei*, 18, var. *molesta*, 19, 76, var. *oloken-ensis*, 20, var. *pteropoda*, 17, var. *wahiauana*, 19, var. *waimeana*, 17, var. *Wilkesii*, 21; *Remyi*, 15, var. *a*, 15, var. *β*, 21; *rivularis*, 33; *Rockii*, 13; *Skottsbergii*, 54, var. *audens*, 55, var. *kalaeloana*, 55, 92, var. *Vaccinioides*, 56; *sparsiflora*, 51; *Stokesii*, 26; *Thymifolia*, 67
- Eurhizocarpon*, 551
- Eurya* subgen. *Euryodes* and *Penteurya*, 299
- Eurya*, 299, 306; *acuminata*, 321, var. *Groffii*, 325; *acuminata* var. *euprista*, 322, var. *monticola*, 323, var. *multiflora*, 322, 325; *acuminatissima*, 328; *albiflora*, 346; *albo-punctata*, 353; *amplexicaulis*, 338; *amplexifolia*, 306; *anceps*, 310; *angulosa*, 353; *angustifolia*, 321, 322; *arbutifolia*, 353; *arisanensis*, 323; *aurescens*, 309; *auriculata*, 338; *Benthamiana*, 353; *bifaria*, 321; *Blumeana*, 321; *boliviensis*, 354; *Bolleana*, 354; *boninensis*, 351; *buxifolia*, 338; *calophylla*, 354; *candicans*, 354; *castaneifolia*, 351; *Cavaleriei*, 354; *cerasifolia*, 326; *cer-
nua*, 354; *ceylanica*, 320; *chinensis*, 319; *chinensis*, 316; *ciliata*, 331; *clandestina*, 322, var. *minor*, 322; *conocarpa*, 342; *confinis*, 322, var. *fusca*, 322; *conocarpa*, 354; *cordata*, 354; *coriacea*, 340; *crenatifolia*, 332; *crispa*, 335; *cuneata*, 317, var. *glabra*, 318; *disticha*, 327; *distichophylla*, 330, var. *Henryi*, 330; *Dombeyana*, 354; *Ekmani*, 354; *emarginata*, 316, var. *microphylla*, 317; *Esquirolii*, 354, 389; *euprista*, 321, var. *heteroidea*, 322; *Fangii*, 318; *fasciculata*, 310; *ferruginea*, 354; *flava*, 338; *Friedrichsthaliana*, 354; *glaberrima*, 334; *glaberrima* var. *acuminata*, 323; *glabra*, 342; *glandulosa*, 315; *gnaphalocarpa*, 332; *grandis*, 351; *Greenmaniana*, 344; *Grisebachii*, 354; *Groffii*, 325; *guatemalensis*, 354; *guianensis*, 354; *Handeliana*, 309; *Hasseltii*, 310; *Hayatai*, 334; *Hellwigii*, 345; *Henryi*, 330; *Hintoni*, 354; *hirsutula*, 322; *hortensis*, 335; *inaequifolia*, 354; *inaequilatera*, 354; *Jacquemartii*, 336; *japonica*, 335, var. *microphylla*, 337; *japonica* var. *acuminata*, 322, var. *aurescens*, 314, var. *chinensis*, 320, var. *conocarpa*, 343, var. *hortensis*, 335, var. *montana*, 335, var. *multiflora*, 336, var. *nitida*, 311, var. *parvifolia*, 319, var. *phyllanthoides*, 322, var. *pumila*, 337, var. *pusilla*, 335, var. *Thunbergii*, 311, 336, 343, var. *uniflora*, 335, var. *yakushimensis*, 353; *Karsteniana*, 354; *ladronica*, 311; *lanata*, 354; *lanceifolia*, 354; *lasiopetala*, 354; *latifolia*, 336; *Lehmannii*, 354; *leptantha*, 349; *leptophylla*, 332; *littoralis*, 316; *longipes*, 355; *Loquaiana*, 326; *lucida*, 321; *Macartneyi*, 306, var. *hainanensis*, 307; *macrophylla*, 355; *Matsudai*, 323; *mezophylla*, 348; *membranacea*, 322; *Merrilliana*, 347; *mexicana*, 355; *microphylla*, 337; *monsonensis*, 355; *montana*, 335; *multiflora*, 321; *muricata*, 352; *myrtifolia*, 310, var. *anceps*, 311, var. *Hasseltii*, 311, var. *polymorpha*, 310, var. *virens*, 311; *myrtilloides*, 355; *Nimanimae*, 355; *nitida*, 310, var. *aurescens*, 314; *nitida*, 355, var. *strigillosa*, 323; *obliquifolia*, 315; *obovata*, 343, var. *sinaboengensis*, 344; *obovata* var. *elliptica*, 343; *oreogena*, 346; *osimensis*, 352; *oxysepala*, 350; *pachyphylla*, 340; *pachyrachis*, 340; *palauensis*, 311; *panamensis*, 829; *parvifolia*,

319; *patentipila*, 331; *phyllanthoides*, 322, var. *brevifolia*, 322; *phyllopoda*, 350; *Pickeringii*, 340; *polyneura*, 307; *ponapensis*, 311; *pumila*, 337; *pusilla*, 336; *rapensis*, 352; *Rehderiana*, 349; *rengchiensis*, 337; *reticulata*, 343, 355, var. *subintegrifolia*, 355; *Riehii*, 341; *Roemeri*, 344; *roraimensis*, 355; *rostrata*, 322; *Roxburghii*, 310; *ryukyuensis*, 353; *salicifolia*, 322; *sanguinea*, 341; *Seemanniana*, 355; *sericea*, 355, var. *chrysophylla*, 355; *serrata*, 322, var. *firma*, 322, var. *membranacea*, 322; *spathulifolia*, 355; *stenophylla*, 308; *strigillosa*, 331; *suberosa*, 355; *subintegrifolia*, 355; *Suzukii*, 323; *Swinglei*, 330; *symplocina*, 326; *syphilitica*, 355; *systyla*, 311; *ternatana*, 353; *ternstroemioides*, 355; *tetragonoelada*, 308; *tigang*, 347; *titang* var. *meisophylla*, 348, var. *phyllopoda*, 350; *timorensis*, 353; *trichocarpa*, 328; *trichogyna*, 328; *tristyla*, 342; *uniflora*, 335; *vaccinioides*, 355; *velutina*, 329; *verrucosa*, 355; *virens*, 311, var. *elliptica*, 311; *vitiensis*, 341; *Wallichiana*, 322, 326; *Wawrai*, 355; *Weissiae*, 315; *Wightiana*, 310; *Wrayi*, 323; *yaeyamensis*, 333; *yakushimensis*, 353; *Yungasiae*, 355; *zigzag*, 334; *Zollingeri*, 310

Euryodes (subgen.), 299, 300, 301, 306

Euryptera, 228; *Hassei*, 236; *Howellii*, 237; *insularis*, 239; *luoida*, 235; *palida*, 239; *parvifolia*, 238

F

Ferula caruifolia, 252; *farinosa*, 242; *foeniculacea*, 259; *macrocarpa*, 264; *nudicaulis*, 290; *Nuttallii*, 291; *parvifolia*, 238; *pubescens*, 256

Frangipani, 204

Frangipanier, 204

Freziera, 299, 300; *angulosa*, 353; *arbutifolia*, 353; *boliviensis*, 354; *Bolleana*, 354; *calophylla*, 354; *candicans*, 354; *cernua*, 354; *conocarpa*, 354; *cordata*, 354; *Dombeyana*, 354; *Ekmani*, 354; *ferruginea*, 354; *Friedrichsthaliana*, 354; *Grisebachii*, 354; *guatemalensis*, 354; *guianensis*, 354; *Hieronyma*, 355; *inaequalifolia*, 354; *inaequilatera*, 354; *integrifolia*, 353; *Karsteniana*, 354; *lanata*, 354; *lanceifolia*, 354; *Lehmannii*, 354; *longipes*, 355; *macrophylla*, 355; *mexicana*, 355; *monsonensis*, 355; *Nimanimae*,

355; *reticulata*, 355, var. *subintegrifolia*, 355; *roraimensis*, 355; *Seemanniana*, 355; *sericea*, 355, var. *chrysophylla*, 355; *spathulifolia*, 355; *suberosa*, 355; *subintegrifolia*, 355; *ternstroemioides*, 355; *vaccinioides*, 355; *verrucosa*, 355; *Wawrai*, 355; *Yungasiae*, 355

Fungal components of lichens, 521, 735

G

Gametophyte of *Tradescantia*, granules in, 455

Garland Hershey Ridge, plant life on, 486

Gasparrinia, 622; *cirrochrooides*, 626; *inordinata*, 626; *Joannae*, 626; *lucens*, 626; *Siplei*, 624, 696, 698, 718; *sublobulata*, 627

Geeria angustifolia, 321; *glabra*, 342; *obovata*, 343; *serrata*, 321; *serrata sericea*, 321

Geographical distribution of the Antarctic flora, 467

Geology of Marie Byrd Land, 472

Gloeocapsa, 741, 742, 751

Glossodium, 734

Gomphillus, 734

Gonidia, cephalodial, in *Stereocaulon*, specificity of, 750; variation of, 752

Gonzalagunia panamensis, 837; *rosea*, 836

Graham Land, explorations in, 468, 503, 515

Grant, Adele Lewis. A monograph of the genus *Hemimeris*, 435

Granules in microspores of *Tradescantia*, Optical and chemical studies on the, 455, 466

Graphidineae, 736

Greeneostachys (subsect.), 109

Greenman, Jesse More, portrait of, 1; dedication of number of *ANNALS* in honor of birthday of, 1; Studies of South American *Senecios*—II, 795

Grimmia Antarctica, 719, 721, var. *percompacta*, 723, var. *pilifera*, 722

Grimmiaceae, 721

Guettarda chiriquensis, 838; *crispiflora*, 839

Gymnoderma, 734

Gymnosiphon suaveolens, 825

Gynandra, 299

H

Haines Mountains, geology and plant life of, 486, 487, 506, 508

Hawaiian species of *Euphorbia* L., Revision of the, 1

Hemimeris, A monograph of the genus, 435

Hemimeris, 435, 441; *acutifolia*, 453; *alsinoides*, 446; *bonae-spei*, 435, 436, 452; *caulialata*, 453; *centrodes*, 439, 440, 450; *coccinea*, 453; *diffusa*, 438, 453; *elegans*, 453; *gracilis*, 439, 440, 451, 452; *hirsuta*, 453; *incisifolia*, 453; *intermedia*, 453; *latipes*, 446; *linearifolia*, 453; *macrophylla*, 453; *montana*, 436, 437, 439, 446, 447, var. *pachyceras*, 440, 449; *montana* var. *latipes*, 446; *Mutisi*, 453; *nana*, 450; *pachyceras*, 449; *parviflora*, 453; *peduncularis*, 453; *procumbens*, 453; *sabulosa*, 436, 437, 439, 443, 444; *sessilifolia*, 446; *sinuata*, 446; *unilabata*, 453; *urticifolia*, 453

Heteromyces, 734

Hexanthus umbellatus, 396

Himatanthus Willd., An evaluation of the genera *Plumeria* L. and, 189

Himatanthus, 190, 192; *articulata*, 196, 221, 222, 223; *attenuata*, 197, 222; *bracteata*, 200, 222, 223; *lancifolia*, 200, 222; *obovata*, 201, 222, 223, 224, var. *puberula*, 201, 222, 223, var. *typica*, 201; *phagedaenica*, 199, 221, 222, 223; *rigida*, 193, 196; *Sucuba*, 198, 222, 223

Hormiscium sp., 496, 521, 712

Huea, 617; *cerussata*, 618; *coralligera*, 618; *flava*, 618, 696

Hybridization, evidences of, in the genus *Plumeria*, 217

Hybrids in *Salvia*, 143

I

Ilex crenata luzonica, 355; *emarginata*, 316

Imbricaria, 589

Icososte chinensis, 386, var. *rotundifolia*, 385; *lancifolia*, 406; *pedunculata*, 414; *rotundifolia*, 385, var. *oblongifolia*, 386

Ixora floribunda, 838

J

Jepsonia (subsect.), 118

Johnson, George Thomas. The taxonomic importance and phylogenetic significance of the cephalodia of *Stereocaulon*, 729; Francis O. Schmitt and. Optical and chemical studies on the granules in microspores of *Tradescantia*, 455

Jonker, F. P., Panamanian Burmanniaceae determined by, 825

K

Killip, Ellsworth P., Panamanian Passifloraceae determined by, 830

King Edward VII Land and north-western Marie Byrd Land, map of, 475; lichens of, 500, 515; mosses of, 719

Kobuski, Clarence E. Studies in Theaceae. III. *Eurya* subgenera *Euryodes* and *Penteurya*, 299

Kuttlingeria, 614; *rufa*, 615, 694, 718; *rutilans*, 616, 694, 696

L

Lauraceae, Studies in the. I. Chinese and Indo-Chinese species of *Litsea*, *Neolitsea*, and *Actinodaphne*, 361

Laurus Cubeba, 368; *obovata*, 405; *pi-losa*, 404; *rotundifolia*, 385

Lecania, 583

Lecanora, 568, 736; *carbonacea*, 576, 680; *chrysoleuca* var. *melanophthalma* f. *exsulans*, 570; *cinericola*, 622; *elegans* f. *lucens*, 626; *exsulans*, 570; *fuscobrunnea*, 577, 684; *griseo-marginata*, 572, 682; *inordinata*, 626; *Joannae*, 626; *lilacina*, 574, 682; *lilacinofusca*, 575, 684; *rubina* var. *melanophthalma* f. *exsulans*, 570; subg. *Rinodina*, 655; *Siplei*, 571, 682, 714; *sublobulata*, 627; *subolivacea*, 578, 684

Leibergia, 228; *orogenioides*, 242

Lecanoraceae, 567

Lecidea, 528, 736; *Blackburni*, 540, 670; *Byrdii*, 536, 668; *cancriformis*, 539, 670, 714, *capsulata*, 533, 664; *cerussata*, 618; *coralligera*, 618; *Coreyi*, 534, 668; *ecorticata*, 537, 668; *Painei*, 542, 670; *Siplei*, 530, 666, 714; *Stancikii*, 538, 668, 670; *Wadei*, 532, 666

Lecideaceae, 527

Leptotaenia salmoniflora, 292

Lichen aurantiacus, 622; subg. *Physcia*, 660

Lichen Peak, geology and plant life of, 486, 487

Lichens, 515, 729; algal symbionts, 520, 735; bacterial symbionts, 522; cephalodia of, 729; cortex, 519; fungal symbionts, 521, 735; light relations, 477; lower cortex, 520; medulla, 519; reproduction of, 521; water relations, 485; wind relations, 483; and lichen parasites of the Second Byrd Antarctic Expedition, 467, 497, 515

- Lindera communis*, 399; *Esquirolii*, 369; *fruticosa*, 399; *megaphylla* f. *touyunensis*, 400; *obovata*, 367; *Playfairii*, 400, 431; *populifolia*, 367; *puberula*, 370; *setchunensis*, 411; *subcaudata*, 431
- Lipoids, tests for, in microspore granules of *Tradescantia*, 458
- Litsea*, 362; *acutivena*, 389; *amara*, 396; *aurata*, 398, 422; *auriculata*, 366; *Balansea*, 396; *baviensis*, 376, var. *szemaois*, 377; *baviensis* var. *venulosa*, 378; *brevipetiolata*, 373, 374; *brideliifolia*, 385; *cambodiana*, 383, var. *longiracemosa*, 383, var. *multiracemosa*, 383; *Cavaleriei*, 398; *Chaffanjonii*, 398; *chartacea*, 377, var. *subtriplinervis*, 378; *Chenii*, 372; *chinensis*, 386; *Chunii*, 372; *citrata*, 368; *Clemensii*, 376; *confertifolia*, 399; 416; *consimilis*, 399, 426; *coreana*, 399; *Cubeba*, 368; *cupularis*, 399, 405; *Dunniana*, 391, 431; *Eberhardtii*, 378, var. *subtriplinervis*, 378; *elongata*, 390, var. *cuneifolia*, 391; *Esquirolii*, 389, 399; *eugenioides*, 380; *euosma*, 368; *Faberi*, 390; *Faberi* var. *ganchouensis*, 389; *firma austrannamensis*, 387; *Forrestii*, 369; *fruticosa*, 399; *Garrettii*, 388, var. *longistaminata*, 388; *glauca*, 399, 421; *glutinosa*, 384, var. *brideliifolia*, 385; *gracilipes*, 399, 419; *grandifolia*, 375; *Greenmaniana*, 394; *Griffithii* *annamensis*, 375; *Helferi* *laosensis*, 379; *hexantha*, 396; *honghoensis*, 375; *hupehana*, 391; *hupehana* var. *longifolia*, 399, 413; *ichangensis*, 371; *iteodaphne*, 391, f. *chinensis*, 393; *Kobuskiana*, 354, 389, 431; *lanceifolia*, 395, var. *pedicellata*, 396; *lanceifolia*, 406; *lanceilimba*, 379; *laxiflora*, 399; *Liyuyingii*, 380; *longipes*, 379; *longepetiolata*, 367; *Maclurei*, 376; *Mairei*, 399; *mekongensis*, 392; *mollifolia*, 367; *mollifolia* var. *glabrata*, 369; *mollis*, 367, var. *glabrata*, 369; *monopetala*, 387; *moupinensis*, 370; *multiumbellata*, 373, f. *annamensis*, 373; *myricopsis*, 399; *obovata*, 399, 405; *Panamouja*, 380; *Pierrei*, 382, var. *grandifolia*, 382, var. *lobata*, 382; *Pierrei* var. *szemaois*, 377; *Playfairii*, 400; *polyantha*, 387, 397, f. *glabra*, 398; *populifolia*, 367; *pseudoelongata*, 384; *pulchella*, 400, 427; *pungens*, 371; *Rehderiana*, 381; *rotundifolia*, 385, 414, var. *oblongifolia*, 386, 414; *rubescens*, 372, var. *tonkinensis*, 373, var. *yunnanensis*, 372; *salicifolia*, 397, f. *attenuata*, 397, f. *glabra*, 398; *Sebifera*, 384, var. *brachyphylla*, 385; *sericea*, 369; *shweliensis*, 400, 401; *Thorelii*, 384; *touyunensis*, 400; *umbellata*, 396; *umbrosa*, 400, 426; *undulatifolia*, 400, 417; *Fang*, 382, var. *grandifolia*, 382, var. *lobata*, 382; *variabilis*, 393, var. *oblonga*, 393; *variabilis* var. *tonkinensis*, 393; *Veitchiana*, 370; *verticillata*, 373, f. *annamensis*, 373, var. *brevipes*, 374, var. *brevipetiolata*, 374; *viridis*, 391, var. *Clemensii*, 391; *wenshanensis*, 375; *Wilsonii*, 388; *wushanica*, 400, 419; *zeylanica*, 400, 427, var. *chinensis*, 400, 427
- Little America, 467
- Lobaria amplissima*, 731; *herbaceae*, 731; *linita*, 731; *oregana*, 731
- Lomatia*, 225, 226
- Lomatium*, A revision of the genus, 225
- Lomatium*, 228; *alatum*, 253, 287, var. *purpureum*, 254; *alpinum*, 279; *ambiguum*, 246; *ambiguum* var. *leptocarpum*, 246; *angustatum*, 268, var. *flavum*, 269; *anomalum*, 285; *argense*, 263; *artemisiarum*, 264; *athamantoides*, 259; *Austinae*, 257; *bicolor*, 246; *Brandegei*, 290; *brevifolium*, 250; *Canbyi*, 244; *caruifolium*, 252, var. *denticulatum*, 253, var. *marginatum*, 253, var. *purpureum*, 254; *caruifolium* var. *erythropodium*, 252, var. *solanense*, 252; *Chandleri*, 250; *ciliolatum*, 276; *circumdatum*, 249; *concinnum*, 276; *Congdoni*, 266; *Cous*, 247; *Cusickii*, 287; *dasyacarpum*, 262; *dasyacarpum* var. *decorum*, 262, var. *medium*, 262; *daucifolium*, 225, 260; *Donnellii*, 271; *Eastwoodae*, 275; *ellipticum*, 265; *Engelmannii*, 268; *farinosum*, 242; *flavum*, 264; *foeniculaceum*, 225, 259; *Geyeri*, 243; *giganteum*, 285; *Gormanii*, 240, f. *purpureum*, 241; *Grayi*, 270, var. *depauperatum*, 270; *Greenmanii*, 274; *Hallii*, 279; *Hendersonii*, 244; *Howellii*, 237; *insulare*, 239; *Jonesii*, 258; *juniperinum*, 259; *laevigatum*, 288; *lapidosum*, 292; *latilobum*, 281; *Leibergii*, 279; *Lemmoni*, 292; *leptocarpum*, 245; *lucidum*, 235; *lucidum* var. *repostum*, 237; *MacDougali*, 258; *macrocarpum*, 264, var. *ellipticum*, 265; *macrocarpum* var. *artemisiarum*, 264, var. *Douglasii*, 264, var. *semivittatum*, 264; *marginatum*, 253, var. *purpureum*, 254; *Martindalei*, 280, var.

angustatum, 268; *megarrhizum*, 282; *microcarpum*, 279; *millefolium*, 270; *minimum*, 273; *mohavense*, 263; *montanum*, 248; *Nelsonianum*, 272; *nevadense*, 254, var. *Parishii*, 255, var. *pseudorientalis*, 255; *nevadense* var. *cupulatum*, 254, var. *holopterum*, 255; *nudicaule*, 290; *nudicaule* var. *puberulum*, 285; *Nuttallii*, 278, var. *alpinum*, 279; *oreganum*, 273; *orientale*, 225, 261; *orogenioides*, 242; *Parishii*, 255; *Parryi*, 277; *parvifolium*, 238, var. *pallidum*, 239; *Piperi*, 241; *platycarpum*, 283; *platyphyllum*, 291; *Plummerae*, 256, var. *Austinae*, 257, var. *Helleri*, 257, var. *Sonnei*, 257; *pubescens*, 259; *purpureum*, 248; *repositum*, 237; *rigidum*, 239; *robustius*, 286; *Sandbergii*, 266; *scabrum*, 275; *semisepultum*, 258; *serpentinum*, 271; *simplex*, 282, var. *leptophyllum*, 283; *Sonnei*, 257; *Sukendorffii*, 289, var. *Thompsonii*, 289; *tomentosum*, 263; *Torreii*, 267; *tritermum*, 284, var. *alatum*, 287, var. *anomalum*, 285, var. *brevifolium*, 286, var. *macrocarpum*, 286; *utriculatum*, 250, var. *papillatum*, 251; *utriculatum* var. *anthemifolium*, 250, var. *glabrum*, 250; *vaginatum*, 249; *Vaseyi*, 251; *villosum*, 225, 259; *Watsoni*, 245

Ludvigia alternifolia var. *pubescens*, 772

Lyngbya, 751

M

Machilus hainanensis, 404; *pilosa*, 404; *Thunbergii*, 399

Magnolia guatemalensis, 829; *poasana*, 829; *sorum*, 828; *Yoroconte*, 829

Magnoliaceae, Panamanian, 828

Malapoenna Sieboldii, 421

Marathrum Allenii, 827; *foenicula-ceum*, 828; *Schiedeanum*, 828

Marie Byrd Land: discovery and exploration of, 467; flora of, 468, 486, 497, 501, 719; geology of, 468, 472; history of, 467; lichens and lichen parasites collected in, 515; light in, 477; map of, 475; moisture in, 484; Queen Maud Geological Party in, 471, mosses collected by, 719; rocks in, 486, 490; temperatures in, 478; wind in, 482; summary of plant life of, 497; zoological observations in, 470

Mathias, Mildred E. A revision of the genus *Lomatium*, 225

Maxon, W. R., Panamanian Polypodiaceae determined by, 824

Meristotheca, 299

Merorinis, 655

Microchaete trichopus, 820

Microspores of *Tradescantia*, Optical and chemical studies on the granules in, 455

Missouri: Additions, corrections, and revisions to the Annotated Catalogue of the Flowering Plants of, 775; New varieties and forms from, 769

Moisture in the Antarctic, 484

Monograph of the genus *Hemimeris*, A, 435

Mosses: in Marie Byrd Land, 467, 500; of the second Byrd Antarctic Expedition, 719

Mt. Corey, geology and plant life of, 494, 514

Mt. Donald Woodward, geology and plant life of, 487, 508, 512, 514

Mt. Grace McKinley, geology and plant life of, 476, 491, 506, 508

Mt. Helen Washington, geology and biology of, 481, 486, 490, 501, 504, 506

Mt. Raymond Fosdick, plant life of, 494

Mt. Rea-Cooper, geology and plant life of, 492, 506, 508, 510, 512, 514

Mt. Stancliff, geology and plant life of, 489, 510

Mountains in Marie Byrd Land: granitic, 490; metamorphic sediments and associated dike rocks, 486

Munzia, 100, 104

Musenium tenuifolium, 282

Myrica adenophora, 399; *esculenta*, 399

Myrioblastus, 554

Myrsinaceae, Panamanian, determined by P. C. Standley, 832

Myxophyceae, 520, 751

N

Neea acuminatissima, 823

Neolitsea, 415; *alongensis*, 424; *aurata*, 398, 422, var. *glabrescens*, 422; *cam-bodiana*, 418, var. *glabra*, 418; *chinensis*, 422; *Chunii*, 429, f. *annamensis*, 430; *confertifolia*, 399, 414, 416; *claeocarpa*, 421; *ellipsoidea*, 428; *ferruginea*, 418; *glauca*, 421; *gracilipes*, 419; *homilantha*, 419; *hongkongensis*, 414, 418; *Howii*, 424; *kwangsiensis*, 423; *lanuginosa chinensis*, 422; *Levinii*, 423; *Merrilliana*, 425; *oblongifolia*, 417; *obtusifolia*, 420; *phanerophlebia*, 430, f. *glabra*, 430; *Playfairii*, 400, 431; *Poillanei*,

- 425; polycarpa, 429; pulchella, 400, 427; Sieboldii, 399, 421; subcaudata, 431; subfoveolata, 429; umbrosa, 399, 400, 426; undulatifolia, 400, 417; wushanica, 399, 400, 419; zeylanica, 400, 427, var. Fangii, 428; zeylanica var. obovata, 425
- Neuropogon*, 601; *melaxanthum*, 602; *Taylori*, 602
- New varieties and forms from Missouri, 769
- Nostoc, 739, 740, 741, 742, 744, 751
- Nyctaginaceae, Panamanian, determined by P. C. Standley, 828
- O
- Omphalodium quartum*, 561
- Optical and chemical studies on the granules in microspores of *Tradescantia*, 455
- Orchidaceae, Panamanian, determined by Charles Schweinfurth, 825
- Organ Pipe Mountains, 514
- Orobanchaceae, 835
- Oxypolis* sp., 292; *rigidior*, 292; *ternata*, 292
- P
- Paederota*, 436, 442; *bonae spei*, 436, 437, 438, 443; *racemosa*, 446
- Palicourea panamensis*, 839
- Palmer, Ernest J., and Julian A. Steyermark. New varieties and forms from Missouri, 769; Additions, corrections, and revisions to the "Annotated Catalogue of the Flowering Plants of Missouri," 775
- Panama, Contributions toward a flora of, II. Miscellaneous collections during 1936-1938, 823
- Pannariaceae, 736
- Pannoparmelia*, 586; *delicata*, 587, 714, 716; *pellucida*, 586, 714, 716
- Parishiella* (subsect.), 100, 112
- Parmelia*, 589; *Coreyi*, 595, 688; *griseola*, 596, 688; *leucoblephara*, 592; *quarta*, 561; *variolosa*, 593, 688; parasite of, 662
- Parmeliaceae, 584
- Passiflora salvadorensis*, 830
- Passifloraceae, Panamanian, determined by Ellsworth P. Killip, 830
- Pastinacea ferulacea*, 259; *foeniculacea*, 259; *nudicaulis*, 290
- Peltigera aphthosa*, 757; *cephalodia* of, 730
- Peltigeraceae, 736
- Penicillium* sp., 522
- Penteurya* (subgen.), 300, 301, 345
- Peperomia Alleni*, 825; *cocleana*, 826
- Pepsin, effect of, on microspore granules in *Tradescantia*, 459
- Peucedanum*, 225, 228; *abrotanifolium*, 247; *ambiguum*, 243, 246, var. *leptocarpum*, 246; *argense*, 263; *Austinae*, 257; *bicolor*, 246, var. *gumbonis*, 246; *Brandegeti*, 290; *californicum*, 238, 252; *Canbyi*, 242; *caruifolium*, 252; *circumdatum*, 249; *confusum*, 240; *Cous*, 247; *Cusickii*, 287; *dasyacarpum*, 262; *daucifolium*, 260; *Donnellii*, 271; *erosum*, 253; *euryacarpum*, 264; *Euryptera*, 235; *evittatum*, 243; *farinosum*, 242; *foeniculaceum*, 259, 260, 262, var. *daucifolium*, 259, 260; *Geyeri*, 243; *Gormanii*, 240; *graveolens*, 278, var. *alpinum*, 279; *Grayi*, 270, var. *aberrans*, 270; *Hallii*, 270, 279; *Hassei*, 235; *Hendersonii*, 244; *Howellii*, 237; *insulare*, 239; *Jaredi*, 262; *juniperinum*, 259; *Kingii*, 278, var. *alpinum*, 279; *laevigatum*, 258; *lapidosum*, 292; *latifolium*, 291; *leiocarpum*, 291, var. *campestre*, 291; *Lemmonti*, 292; *leptocarpum*, 245; *ludovicianum*, 292; *macrocarpum*, 264, 277, var. *euryacarpum*, 264; *marginatum*, 253; *Martindalei*, 280, var. *angustatum*, 268; *megarrhiza*, 282; *microcarpum*, 279; *millefolium*, 270; *mohavense*, 263; *montanum*, 248; *nevadense*, 254, var. *cupulatum*, 254; *Newberryi*, 292; *nudicaule*, 254, 261, 279, 291, var. *ellipticum*, 265, var. *puberulum*, 285; *Nuttallii*, 284, 291; *oreganum*, 273; *orientale*, 261; *Parishii*, 255; *Parryi*, 277; *parvifolium*, 238; *Plummerae*, 256; *Pringlei*, 262; *rigidius*, 292; *robustum*, 291; *salmoniflorum*, 292; *Sandbergii*, 266; *scopulorum*, 277; *simplex*, 283; *Suksdorfii*, 289; *tenuissimum*, 247; *ternatum*, 292; *tomentosum*, 263; *Torreyi*, 267; *triternatum*, 245, var. *alatum*, 287, var. *brevifolium*, 286, var. *leptocarpum*, 246, var. *leptophyllum*, 283, var. *macrocarpum*, 286, var. *platycarpum*, 282, var. *robustius*, 286; *utriculatum*, 250; *Vaseyi*, 251; *villosum*, 225, 259, 260; *Watsoni*, 240, 245
- Phalodictyum*, 550
- Phragmothele*, 524; *papulare*, 524
- Phylogenetic significance of the cephalodia of *Stereocaulon*, The taxonomic importance and, 729
- Physcia, 660
- Physcia*, 589; *cirrhochroa*, 620

- Physciaceae, 660
 Pilophoron, 734, 736, 747
 Piper Alleni, 826; *canyazasense*, 826;
 san joseanum var. *chiriquinum*, 826;
tabasaranum, 827; *yapeanum*, 827
 Piperaceae, Panamanian, determined by
 William Trelease, 825
Placodium cirrochrooides, 626; *inordi-*
nata, 626; *Joannae*, 626; *lucens*, 627;
sublobulatum, 627; sect. *Thamnomia*,
 627
Pleolecis, 556
Pleurinis, 655
 Plumeria L.: evidences of hybridiza-
 tion in the genus, 217; and Himatan-
 thus Willd., An evaluation of the
 genera, 189
 Plumeria, 189, 202; *acuminata*, 207;
acutifolia, 208, β . *Gasparrini*, 208;
alba, 216; *alba* var. *fragrans*, 206,
 β . *Jacquiniana*, 216; *ambigua*, 199,
 221; *angustiflora*, 197; *angustifolia*,
 221; *apiculata*, 213; *articulata*, 196,
 221, 222, 223; *attenuata*, 197, 221,
 β . *obtusifolia*, 197, γ . *Malongo*, 197;
barahonensis, 212; *beatensis*, 213;
Berteri, 216; *bracteata*, 200, 222,
 223; *castildensis*, 213; *cayensis*, 213;
clusioides, 213; *cochleata*, 206; *cu-*
bensis, 213; *cuneifolia*, 213; *cuspi-*
data, 222; *dictyophylla*, 213; *discolor*,
 220; *drastica*, 196, 222; *Ekmanii*,
 213; *estrellensis*, 213; *fallax*, 196, 199,
 222; *filifolia*, 215; *floribunda*, 198,
 222, δ . *acutifolia*, 199, γ . *calycina*, 199,
 β . *crassipes*, 199; *Hilariana*, 201, 222;
hypoleuca, 216, β . *angustifolia*, 216;
incarnata β . *Milleri*, 208; *inodora*,
 206, 832; *Jaegeri*, 216; *Lambertiana*,
 208; *lanceifolia*, 200, 222, β . *major*,
 199, 200, γ . *microphylla*, 199; *leuco-*
neura, 214; *longiflora*, 220; *loranthi-*
folia, 208; *Malongo*, 197; *Martii*,
 199, 223; *megaphylla*, 208; *micro-*
calyz, 196, 223; *Milleri*, 208; *mollis*,
 208; *multiflora*, 215; *obovata*, 201,
 223, 224; *obtusata*, 212, 218, var. *serici-*
folia, 214, var. *typica*, 212; *oligo-*
neura, 201, 223; *Ostenfeldii*, 213;
phagedaenica, 199, 221, 222, 223;
pilosula, 214; *puberula*, 201, 222, 223,
 224; *pudica*, 206; *revoluta*, 200, 223;
rubra, 202, 207, 218, f. *acutifolia*,
 211, f. *lutea*, 211, f. *tricolor*, 211,
 f. *typica*, 211; *rubra* \times *subsessilis*,
 220; *sericifolia*, 214; *speciosa*, 199,
 223; *stenopetala* var. *angustissima*,
 220; *stenophylla*, 215; *subsessilis*,
 216, 217; *Succuba*, 198, 222, 223;
tarapotensis, 198, 223; *Tenorii*, 212;
trouinensis, 220; *tuberculata*, 214;
velutina, 201, 223; *Warmingii*, 201,
 224
Plumiera, 202; *alba* var. *fragrantissima*,
 206, var. *inodora*, 206; *arborescens*,
 208; *caracasana*, 206; *confusa*, 212;
inaguensis, 212; *jamaicensis*, 212;
Kerrii, 208; *lanata*, 214; *latifolia*,
 222; *montana*, 213; *nipensis*, 212;
trinitensis, 214
Plumieria, 203; *bahamensis*, 212; *bi-*
color, 207; *biglandulosa*, 220; *car-*
inata, 207; *clusioides*, var. *parviflora*,
 212; *domingensis*, 214; *emarginata*,
 212, β . *sericifolia*, 214; *gibbosa*, 214;
incarnata, 207; *Jamesoni*, 208;
Krugii, 212; *latifolia*, 201; *lutea*,
 208; *Marchii*, 212; *mexicana*, 208;
obtusata γ . *laevis*, 212, β . *parviflora*,
 212; *Paulinae*, 220; *portoricensis*,
 212; *purpurea*, 207; *sericifolia*, 214;
tricolor, 207; *venosa*, 212
 Podostemonaceae, Panamanian, 827
Poinsettia (sect.), 70
Poinsettia, 2, 70; *cyathophora*, 72; *Ed-*
wardsii, 70; *heterophylla*, 70
 Polycauliona, 627; *pulvinata*, 628, 696,
 718; *regalis*, 627; *sparsa*, 629, 696
 Polypodiaceae, Panamanian, determined
 by William R. Maxon, 824
Proteurya, 299, 300, 306
Protoblastenia, 605; *alba*, 607, 692;
aurea, 608, 692; *citrinigracans*, 610,
 692; *flava*, 605, 692
 Proteins, tests for, in microspore gran-
 ules of Tradescantia, 458
Pseudocymopterus humboldtensis, 292;
montanus, 292
Pseudocurya crenatifolia, 332
Pycnosphace (subsect.), 107
Pycnosphace (sect.), 107
Pycnosphace, 107; *Columbariae*, 107
 Pyrenocarpeae, 736
 Pyrenodesmia, 619; *athallina*, 621; *au-*
rantia, 622; *climicola*, 622; *Dar-*
bishirei, 620, 712, 714
- Q
- Queen Maud Mountains, geology and
 plant life of, 467, 495, 514; lichens of,
 515; temperature of, 480
Quercus falcata f. *angustior*, 771
- R
- Ramona*, 98; *capitata*, 123; *Clevelandii*,
 119; *Dorrii*, 133; *grandiflora*, 110;
humilis, 136; *incana*, 130; *nivea*, 121;
pachystachya, 125; *Palmeri*, 146;

- pilosa*, 133; *polystachya*, 140; *stachyoides*, 113; *Vaseyi*, 138
Rapanea aurea, 311
Ravnia triflora, 838
 Raymond Fosdick Mts., geology and plant life of, 494
Rehmiopsis, 551
 Revisions, Additions, corrections, and, to the "Annotated Catalogue of the Flowering Plants of Missouri," 775
 Revision of the genus *Lomatium*, A, 225
 Revision of the Hawaiian species of *Euphorbia* L., 1
Rhizocarpon, 550; *flavum*, 552, 674, 676, 718
Rinodina, 655, 712; *olivaceobrunnea*, 659, 710; *sordida*, 657, 710
 Rocks in Marie Byrd Land: granitic, 473, 490; metamorphic sedimentary, 486
 Rubiaceae, Panamanian, determined by P. C. Standley, 836
- S
- Salix cordata* f. *mollis*, 770, f. *subintegra*, 770; sericea f. *glabra*, 769
Salvia, 95; *apiana*, 140, 186, 188; *apiana* var. *compacta*, 141, var. *typica*, 140; *apiana* × *Clevelandii*, 146; *apiana* × *eremostachya*, 148; *apiana* × *leucophylla*, 147; *apiana* × *mellifera*, 146; *apiana* × *Munzii*, 148; *apiana* × *pachyphylla*, 147; *apiana* × *Vaseyi*, 148; *bernardina*, 145, 149; *Brandegei*, 117, 168; *californica*, 104, 154; *californica*, 140; *carduacea*, 101, 152; *carnosa*, 128, subsp. *argentea*, 131, subsp. *Gilmani*, 132, subsp. *Mearnsii*, 131, subsp. *pilosa*, 133, 178, subsp. *typica*, 130; *carnosa* var. *compacta*, 125, var. *pilosa*, 133, var. *typica*, 130; *Clevelandii*, 119, 170; *Clevelandii* × *leucophylla*, 150; *Columbariae*, 107, 145, 160; *Columbariae* × *mellifera*, 145, 149; *Columbariae* var. *bernardina*, 149; *compacta*, 125; *eremostachya*, 134, 180; *eremostachya* × *Vaseyi*, 150; *funerea*, 105, 156; *funerea* var. *formicis*, 105; *gossypina*, 101; *Greatae*, 106, 158; *leucophylla*, 121, 172; *leucophylla* × *mellifera*, 149; *mellifera*, 112, 145, 164; *mellifera* var. *Jonesii*, 152, var. *revoluta*, 117, var. *typica*, 113; *mohavensis*, 123, 174; *Munzii*, 115, 166; *pachyphylla*, 125, 176; *Palmeri*, 146; *pilosa*, 133; *sonomensis*, 136, 182; *spathacea*, 109, 162; *Vaseyi*, 138, 184
- Salvias*, The Californian. A review of *Salvia*, section *Audibertia*, 95
Sambucus canadensis f. *rubra*, 773
Sarcogyne, 556; *angulosa*, 556, 680; *grisea*, 558, 680
Sarconeurum antarcticum, 719; *glaciale*, 719
Sassafras Tzumu, 399
Saunders Mts., 486; geology and plant life of, 493
 Schmitt, Francis O. and G. T. Johnson. Optical and chemical studies on the granules in microspores of *Tradescantia*, 455
 Schweinfurth, Charles, Panamanian Orchidaceae determined by, 825
Scoparia dulcis, 436
Scopulariopsis, *brevicaulis*, 522, 712
 Seudder Mt., plant life of, 496, 512
Scytonema, 740, 742, 751
Sebifera glutinosa, 384
 Second Byrd Antarctic Expedition—Botany, 467
 Seibert, Russell J., R. E. Woodson Jr. and. Contributions toward a flora of Panama. II. Miscellaneous collections during 1936-1938, 823
 Senecio, Studies of South American—II, 795
Senecio, 795; *aberrans*, 796, 806; *abietinus*, 796; *adenotrichus*, 796; *algens*, 796; *algens* var. *major*, 796, var. *minor*, 796; *Andrei*, 797; *apiculatus*, 797; *arbores*, 798, 801; *arbutifolius*, 798; *argenteus*, 798; *Aschenbornianus*, 798; *attenuatus*, 799; *baccharidiflorus*, 799; *bahioides*, 799, a *lanosus*, 799, β *glaber*, 799; *bogotensis*, 799; *Bowmani*, 799; *brachyodon*, 800; *Brittonianus*, 800, 810; *canabinæfolius*, 800; *chachapoyensis*, 801; *chaquiroensis*, 801; *coroicensis*, 806; *dictyophlebius*, 797, 801; *disciformis*, 805; *ellipticifolius*, 802; *ellipticus*, 803; *eristhalifolius*, 803; *erosus*, 816; *evacoides*, 803; *Georgianus*, 803; *grandifolius*, 804; *Greenmanianus*, 803; *Gürkei*, 804; *Hallii*, 803; *Hilli*, 804; *Hypsobates*, 805; *Klattii*, 805, 816; *Klugii*, 806; *laricifolius*, 807; *ledifolius*, 807; *ledifolius* β *Schlimii*, 807; *Lindenii*, 807; *Macbridei*, 808; *magellanicus*, 809; *Magnusii*, 809; *melanolepis*, 809; *Millei*, 809; *modestus*, 810; *modestus*, 822; *morrensis*, 807; *myriocephalus*, 800; *nevadensis*, 810; *octophyllus*, 811; *olophyllus*, 811; *pachypus*, 811; *Pampae*, 812; *pellucidinervis*, 800, 813; *penalis*,

- 813; peregrinus, 800; pericaulis, 814; *pichinchensis*, 814; *pimpinellaefolius*, 815; *polyphyllus*, 815; *praeruptorum*, 816; *pulchellus*, 816; *pulchellus* β . *trichopus*, 820; *rhizomatus*, 816; *Richii*, 816, var. *latior*, 817; *Richii* var. β , 817; *roseus*, 805; *scaphiformis*, 817; *semidentatus*, 808; *Sepium*, 818; *sinuatilobus*, 818; *sotariensis*, 818; *Sprucei*, 800; *stigophlebius*, 819; *subcandidus*, 819; *subdecurrens*, 819; *subglomerosus*, 819; *tephrosioides*, 819; *teretifolius*, 819; *thaeifolius*, 801, 810; *tolimensis*, 820; *trichopus*, 820; *tristis* var. *Comberi*, 820; *Urbani*, 821; *vaccinioides*, 821, var. *pruinosa*, 821; *vernicosus* var. *microphyllus*, 807; *verticillatus*, 821, var. *trichophorus*, 822; *volubilis*, 801; *Werdermannii*, 822; *yungacensis*, 806; *yurensis*, 822
- Seseli biternatum*, 284; *leiocarpum*, 291; *Nuttallii*, 278; *triternatum*, 284
- Sherff, Earl Edward. Revision of the Hawaiian species of *Euphorbia* L., 1
- Siegertia*, 550
- Siple, Paul A. The second Byrd Antarctic Expedition—Botany. I. Ecology and geographical distribution, 467
- Sirosiphon*, 751
- Skua Gull Peak, 481, 486, 488, 510, 512; geology and plant life of, 488
- Skua gulls, 481, 488, 502, 522
- Smyrnium nudicaule*, 290
- Snowy petrel on Mt. Helen Washington, 481, 486, 490, 504, 506
- Solorina, 736; crocea, 736; octospora, 731
- Solubility properties of microspore granules of *Tradescantia*, 459
- South Victoria Land, 468; lichens of, 515
- Sphaerophorus*, 736
- Standley, P. C., Panamanian plants determined by: *Begoniaceae*, 830; *Combretaceae*, 831; *Myrsinaceae*, 832; *Nyctaginaceae*, 828; *Rubiaceae*, 836; *Theaceae*, 829
- Stemmadenia* Donnell-Smithii, 833
- Stereocaulon*, The taxonomic importance and phylogenetic significance of the cephalodia of, 729; phylogenetic connections within the genus, 734, 735
- Stereocaulon*, 729; *albicans*, 746, 748; *alpinum*, 746, 748, 752; *arbuscula*, 746, 748; *botryophorum*, 732, 746, 748; *chlorocarpoides*, 746, 749; *condensatum*, 746, 747, 748; *congestum*, 746, 748; *coralloides*, 746, 748; *cornutum*, 746, 748; *corticatum*, 746, 748; *curtatum*, 746, 748; *denudatum*, 732, 746, 748, 751, 752; *exutum*, 739, 746, 748, 752; *fastigiatum*, 751; *foliolosum*, 746, 748; *glareosum*, 746, 748; *gracilescens*, 746, 748; *grande*, 751; *japonicum*, 739, 746, 748, 752; *lecanoreum*, 746, 749; *macrocarpum*, 746, 749; *macrocephalum*, 746, 749; *Masartianum*, 753; *mixtum*, 746, 749, 752, 753; *nabewaziense*, 746, 748; *nanum*, 746, 748; *nesacum*, 739, 746, 749, 752, 753; *nigrum*, 746, 748; *octomerellum*, 746, 748; *paschale*, 732, 739, 741, 748, 751, 752; *pileatum*, 746, 747, 748; *pilophoroides*, 746, 749; *piluliferum*, 746, 749; *prostratum*, 746, 748, 752; *proximum*, 746, 749, 752, 753; *pygmaeum*, 745; *ramulosum*, 732, 739, 746, 749, 751, 752, 753; *rivalorum*, 746, 748; *sinense*, 746, 749; *sorediiferum*, 746, 749; *sphaerophoroides*, 746, 748; *strictum*, 746, 749; *tomentosum*, 739, 741, 746, 748, 752; *uvuliferum*, 746, 748; *Wrightii*, 746, 748
- Stereopeltis*, 556
- Steyermark, Julian A., Ernest J. Palmer and. New varieties and forms from Missouri, 769; Additions, corrections, and revisions to the "Annotated Catalogue of the Flowering Plants of Missouri," 775
- Stictaceae*, 520, 736
- Stigonema*, 740, 741, 742, 751
- Strangospora*, 554
- Studies in the Apocynaceae. VII. An evaluation of the genera *Plumeria* L. and *Himatanthus* Willd., 189
- Studies in the Lauraceae. I. Chinese and Indo-Chinese species of *Litsea*, *Neolitsea*, and *Actinodaphne*, 361
- Studies in Theaceae. III. *Eurya* subgenera *Euryodes* and *Penteurya*, 299
- Sunlight conditions in Marie Byrd Land, 477
- Symplocarpon* Hintoni, 354
- Symplocos laurina*, 254; *stellaris*, 398
- T
- Taxodium distichum* f. *confusum*, 769
- Taxonomic importance and phylogenetic significance of the cephalodia of *Stereocaulon*, The, 729
- Temperatures: effect of, on pollen granules of *Tradescantia*, 461; in the Antarctic, 478
- Ternstroemia*, 355; *albo-punctata*, 353; *bifaria*, 321

Ternstroemiopsis (subgenus), 299, 300, 301

Tetradenia aurata, 422; *brideliifolia*, 385; *consimilis*, 426; *glauca*, 421; *obovata*, 405; *umbrosa*, 426; *zeylanica*, 427

Tetranthera amara, 396; *attenuata*, 397; *chartacea*, 378; *elongata*, 390; *feruginea*, 396; *glauca attenuata*, 397; *iteodaphne*, 391; *lanceifolia*, 395; *lancifolia*, 395; *macrophylla*, 387; *monopetala*, 387; *obovata*, 405; *Panamaja*, 381; *Panamonja*, 381; *pilosa*, 404; *polyantha*, 387; *rotundifolia*, 385; *salicifolia*, 397; *sericea*, 369; *umbrosa*, 426

Theaceae, Studies in. III. *Eurya* subgenera *Euryodes* and *Penteurya*, 299; Panamanian, determined by P. C. Standley, 829

Thelidium, 524; *Caloplacae*, 525, 664; *inaequale*, 524, 664; *parvum*, 526, 712

Thorne Glacier, 496

Thysanothecium, 734

Tithymalus, 2, 73

Tithymalus (sect.), 73

Tortulaceae, 719

Tradescantia, Optical and chemical studies on the granules in microspores of, 455, 466

Tradescantia paludosa, microspores of, 456; photomicrographs of, 466

Trelease, William, Panamanian Piperaceae determined by, 825

Trypsin, effect of, on microspore granules of Tradescantia, 459

U

Umbilicaria, 559; *cerebriformis*, 562, 678, 714; *cristata*, 565, 678; *pateriformis*, 564, 676; *rugosa*, 561, 676, 714; *spongiosa*, 566, 678

Umbilicariaceae, 559

Urea, reaction of, on pollen granules of Tradescantia, 460

Usnea, 600; *antaretica*, 602, 690; *frigida*, 603, 690; sect. *Neuropogon*, 601; subgen. *Neuropogon*, 601; *sulphurea* f. *sphacelata*, 602

Usneaceae, 597

V

Veronica Africana, 436, 443

Verrucariaceae, 523

Vincetoxicum discolor, 833, 834

Volcano, plant life of, 494

W

Wind in the Antaretics, effect of, on plant life, 482, 502

Woodson, Robert E. Jr. Studies in the Apocynaceae. VII. An evaluation of the genera *Plumeria* L. and *Himantanthus* Willd., 189; and Russell J. Seibert. Contributions toward a flora of Panama. II. Miscellaneous collections during 1936-1938, 823

X

Xanthoria antarctica, 625; *lychnea* f. *antarctica*, 625